


AMENDMENT TO THE CLAIMS

Please amend the claims without prejudice, without admission, without surrender of subject matter, and without any intention of creating any estoppel as to equivalents, as follows.

In the Claims:

1-2. (Cancelled)

3. (Previously presented) The herbicide combination as claimed in claim 19, which comprises, as component (A), one or more triazine derivatives of the formula (X)



(X)

in which

R¹ is (C₁-C₄)-alkyl or (C₁-C₄)-haloalkyl;

R² is (C₁-C₄)-alkyl, (C₃-C₆)-cycloalkyl or (C₃-C₆)-cycloalkyl-(C₁-C₄)-alkyl and

A is -CH₂-, -CH₂-CH₂-, or -CH₂-CH₂-CH₂-.

4-5. (Cancelled)

6. (Previously presented) The herbicide combination as claimed in claim 19, which comprises one or more further components selected from the group consisting of crop protection agents of a different type, additives which are customary in crop protection and formulation auxiliaries.

7. (Previously presented) A method for controlling harmful plants, which comprises applying the herbicides of the herbicide combination as defined in claim 19 together or

separately, pre-emergence, post-emergence or pre- and post-emergence, to the plants, parts of plants, plant seeds or the area under cultivation.

8. (Original) The method as claimed in claim 7 for the selective control of harmful plants in crops of plants.

9. (Previously presented) The method as claimed in claim 7 for the control of harmful plants in cereal.

10-11. (Cancelled)

12. (Previously presented) The herbicide combination as claimed in claim 19, wherein the components are present in a weight ratio (A) : (B) from 1:800 to 3000:1.

13. (Previously presented) The herbicide combination as claimed in claim 2 in which

- halogen is chlorine, bromine or iodine;
- the haloalkyl group is fluoroalkyl
- R¹ is -CF(CH₃)₂,
- R² is (C₁-C₄)-alkyl or (C₃-C₄)-cycloalkyl.
- A is -CH₂-, -CH₂-CH₂- or -CH₂-CH₂-CH₂-.

14. (Previously presented) The herbicide combination as claimed in claim 19, wherein the one or more herbicidally active aminotriazine compound (A) is selected from the group consisting of:

- (A1) 4-amino-6-(1-fluoro-1-methylethyl)-2-(3-phenyl-1-cyclobutylamino)-1,3,5-triazine;
 - (A2) 4-amino-6-(1-fluoro-1-methylethyl)-2-(3-phenyl-1-cyclopropylamino)-1,3,5-triazine;
 - (A3) 4-amino-6-(1-fluoro-1-methylethyl)-2-(3-phenyl-1-ethylpropylamino)-1,3,5-triazine;
- and mixtures thereof; and

wherein the one or more herbicides (B) is selected from the group consisting of:

(B1.1.1) isoproturon;
(B1.2.1) flufenacet;
(B1.2.2) pendimethalin;
fenoxaprop-P-ethyl;
(B1.6.4) MY 100;
(B1.6.5) anilofos;
(B1.6.14) AEB391;
(B2.1.1) tribenuron-methyl;
(B2.1.4) amidosulfuron;
(B2.1.9) ethoxysulfuron;
(B2.2.4) mecoprop-(P);
(B2.3.1) bromoxynil;
(B2.3.2) ioxynil;
(B2.5.2) florasulam;
(B2.6.3) carfentrazone-ethyl;
(B2.6.8) cinidon-ethyl;
(B3.1.4) iodosulfuron-methyl;
(B3.1.5) AEF060;
(B3.1.11) AEF360;
(B3.4.2) diflufenican;
(B3.4.4) flurtamone;
(B3.4.5) isoxaflutole;
(B4.1.2) glufosinate monoammonium salt; and
mixtures thereof.

15. (Previously presented) The herbicide combination as claimed in claim 14,
wherein the one or more herbicidally active aminotriazine compound (A) is:

(A1) 4-amino-6-(1-fluoro-1-methylethyl)-2-(3-phenyl-1-cyclobutylamino)-1,3,5-triazine; and
the one or more herbicides (B) is selected from the group consisting of:

(B1.1.1) isoproturon;
(B1.2.1) flufenacet;

(B1.2.2) pendimethalin;
(B1.3.3) fenoxaprop-P-ethyl and fenoxaprop-ethyl;
(B1.6.4) MY 100;
(B1.6.5) anilofos;
(B1.6.14) AEB391;
(B2.1.1) tribenuron-methyl;
(B2.1.4) amidosulfuron;
(B2.1.9) ethoxysulfuron;
(B2.2.4) mecoprop-(P);
(B2.3.1) bromoxynil;
(B2.3.2) ioxynil;
(B2.5.2) florasulam;
(B2.6.3) carfentrazone-ethyl;
(B2.6.8) cinidon-ethyl;
(B3.1.4) iodosulfuron-methyl;
(B3.1.11) AEF360;
(B3.4.2) diflufenican;
(B3.4.4) flurtamone;
(B3.4.5) isoxaflutole;
(B4.1.2) glufosinate monoammonium salt; and a
mixture of (3.1.4) iodosulfuron-methyl and (B3.1.11) AEF360.

or

the one or more herbicidally active aminotriazine compound (A) is:

(A2) 4-amino-6-(1-fluoro-1-methylethyl)-2-(3-phenyl-1-cyclopropylamino)-1,3,5-triazine; and
the one or more herbicides (B) is:

(B1.1.1) isoproturon;

or

the one or more herbicidally active aminotriazine compound (A) is:

(A3) 4-amino-6-(1-fluoro-1-methylethyl)-2-(3-phenyl-1-ethylpropylamino)-1,3,5-triazine; and

the one or more herbicides (B) is:

(B3.1.4) iodosulfuron-methyl;

(B3.1.5) AEF060; or

a mixture of (B1.1.1) isoproturon and fenoxaprop-P-ethyl;

or

the one or more herbicidally active aminotriazine compound (A) is:

(A4) 4-amino-6-(1-fluoro-1-methylethyl)-2-(2-(3-chlorophenoxy)-1-methylethylamino)-1,3,5-triazine; and

the one or more herbicides (B) is:

(B1.1.1) isoproturon;

(B3.1.4) iodosulfuron-methyl;

(B3.1.5) AEF060; or

a mixture of (B3.1.4) iodosulfuron-methyl and (B3.1.5) AEF060;

or

the one or more herbicidally active aminotriazine compound (A) is:

(A5) 4-amino-6-(1-fluoro-1-methylethyl)-2-(2-(3-chloro-5-methoxyphenoxy)-1-methylethylamino)-1,3,5-triazine; and

the one or more herbicides (B) is:

(B3.1.5) AEF060.

16. (Previously presented) The herbicide combination as claimed in claim 14, wherein the one or more herbicides (B) is selected from the group consisting of:

(B1.3.1) clodinafop-propargyl;

(B1.3.2) diclofop-methyl;

(B1.3.3) fenoxaprop-P-ethyl and fenoxaprop-ethyl;

- (B1.3.4) quizalofop-P and its salts and esters and quizalofop and its salts and esters;
 - (B1.3.5) fluazifop-P and its esters and fluazifop and its esters;
 - (B1.3.6) haloxyfop and haloxyfop-P and their esters;
 - (B1.3.7) propaquizafop;
 - (B1.3.8) cyhalofop and its esters;
 - (B2.2.3) dichlorprop;
 - (B2.2.4) mecoprop-(P); and
- mixtures thereof.

17. (Previously presented) The herbicide combination as claimed in claim 16, wherein the one or more herbicidally active aminotriazine compound (A) is:


(A1) 4-amino-6-(1-fluoro-1-methylethyl)-2-(3-phenyl-1-cyclobutylamino)-1,3,5-triazine; and the one or more herbicides (B) is selected from the group consisting of:

- (B1.3.1) clodinafop-propargyl;
 - (B1.3.2) diclofop-methyl;
 - (B1.3.3) fenoxaprop-P-ethyl and fenoxaprop-ethyl;
 - (B1.3.4) quizalofop-P and its salts and esters and quizalofop and its salts and esters;
 - (B1.3.5) fluazifop-P and its esters and fluazifop and its esters;
 - (B1.3.6) haloxyfop and haloxyfop-P and their esters;
 - (B1.3.7) propaquizafop;
 - (B1.3.8) cyhalofop and its esters;
 - (B2.2.3) dichlorprop;
 - (B2.2.4) mecoprop-(P); and
- mixtures thereof.

18. (Previously presented) The herbicide combination as claimed in claim 17, wherein the one or more herbicidally active aminotriazine compound (A) is:

(A1) 4-amino-6-(1-fluoro-1-methylethyl)-2-(3-phenyl-1-cyclobutylamino)-1,3,5-triazine; and the one or more herbicides (B) is fenoxaprop-P-ethyl.

19. (Previously presented) A herbicide combination comprising a synergistically effective amount of components (A) and (B), wherein component (A) comprises a compound of the formula (II), (III), (IV) and (V), where the compounds are
- compounds of the formula (II) and their salts



in which

R^1 is (C_1-C_6) -alkyl,

which is unsubstituted or substituted by one or more radicals selected from the group consisting of halogen, cyano, nitro, thiocyanato, (C_1-C_4) -alkoxy, (C_1-C_4) -alkylthio, (C_1-C_4) -alkylsulfinyl, (C_1-C_4) -alkylsulfonyl, (C_2-C_4) -alkenyl, (C_2-C_4) -alkynyl, phenyl, which is unsubstituted or substituted, and heterocyclyl having 3 to 6 ring atoms and 1 to 3 hetero ring atoms selected from the group consisting of N, O and S, the ring being unsubstituted or substituted,

R^2 and R^3 in each case independently of one another are hydrogen, amino or alkylamino or dialkylamino having in each case 1 to 6 carbon atoms in the alkyl radical, an acyclic or cyclic hydrocarbon radical or hydrocarbonoxy radical having in each case 1 to 10 carbon atoms or a heterocyclyl radical, heterocycliloxy radical or heterocyclylamino radical having in each case 3 to 6 ring atoms and 1 to 3 hetero ring atoms selected from the group consisting of N, O and S, where each of the five last-mentioned radicals is unsubstituted or substituted, or an acyl radical or


R^2 and R^3 together with the nitrogen atom of the group NR^2R^3 are a heterocyclic radical having 3 to 6 ring atoms and 1 to 4 hetero ring atoms, where the further hetero ring atoms which are optionally present in addition to the nitrogen atom are selected from the group consisting of N, O and S and the radical is unsubstituted or substituted,

R^4 is hydrogen, amino, alkylamino or dialkylamino having in each case 1 to 6 carbon atoms in the alkyl radical, an acyclic or cyclic hydrocarbon radical or hydrocarbonoxy radical

- having in each case 1 to 10 carbon atoms, or a heterocyclyl radical, heterocyclyloxy radical or heterocyclylamino radical having in each case 3 to 6 ring atoms and 1 to 3 hetero ring atoms selected from the group consisting of N, O and S, where each of the five last-mentioned radicals is unsubstituted or substituted, or an acyl radical,
- R^5 is hydrogen, halogen, nitro, cyano, thiocyanato or a radical of the formula $-B^1-Y^1$, where B^1 and Y^1 are as defined below,
- A is an alkylene radical having 1 to 5 straight-chain carbon atoms or alkenylene or alkynylene having in each case 2 to 5 straight-chain carbon atoms, where each of the three last-mentioned diradicals is unsubstituted or substituted by one or more radicals selected from the group consisting of halogen, nitro, cyano, thiocyanato and a radical of the formula $-B^2-Y^2$,
- $(X)_n$ are n substituents X, where X in each case independently of one another, is halogen, (C_1-C_6) -alkyl, (C_2-C_6) -alkenyl, (C_2-C_6) -alkynyl, (C_1-C_6) -alkoxy, (C_2-C_6) -alkenylloxy, (C_2-C_6) -alkynylloxy, $[(C_1-C_4)$ -alkyl]-carbonyl, $[(C_1-C_4)$ -alkoxy]-carbonyl or $[(C_1-C_4)$ -alkylthio]-carbonyl, where the hydrocarbon-containing moieties in the 9 last-mentioned radicals are unsubstituted or substituted, or is a radical of the formula $-B^0-R^0$, where B^0 is as defined below and R^0 is an aromatic, saturated or partially saturated carbocyclic or heterocyclic radical, where the cyclic radical is substituted or unsubstituted, or two adjacent radicals X together are a fused-on cycle having 4 to 6 ring atoms which is carbocyclic or contains hetero ring atoms selected from the group consisting of O, S and N and which is unsubstituted or substituted by one or more radicals selected from the group consisting of halogen, (C_1-C_4) -alkyl and oxo,
- n is 0, 1, 2, 3, 4 or 5,
- B^0, B^1, B^2 in each case independently of one another are a direct bond or a divalent group of the formula $-O-$, $-S(O)_p-$, $-S(O)_p-O-$, $-O-S(O)_p-$, $-CO-$, $-O-CO-$, $-CO-O-$, $-NR^1-$, $-O-NR^1-$, $-NR^1-O-$, $-NR^1-CO-$, $-CO-NR^1-$, where $p = 0, 1$ or 2 and R^1 is hydrogen, alkyl having 1 to 6 carbon atoms, phenyl, benzyl, cycloalkyl having 3 to 6 carbon atoms or alkanoyl having 1 to 6 carbon atoms,
- Y^1, Y^2 in each case independently of one another are H or an acyclic hydrocarbon radical having, for example, 1 to 20 carbon atoms or a cyclic hydrocarbon radical having 3 to 8 carbon atoms or a heterocyclic radical having 3 to 9 ring atoms and 1 to 3 hetero ring atoms selected

from the group consisting of N, O and S, where each of the three last-mentioned radicals is unsubstituted or substituted;

- compounds of the formula (III) or their salts



in which


R^1 is aryl, which is unsubstituted or substituted, or (C₃-C₉)-cycloalkyl, which is unsubstituted or substituted, or heterocyclyl, which is substituted or unsubstituted, or (C₁-C₆)-alkyl, (C₂-C₆)-alkenyl or (C₂-C₆)-alkynyl,

where each of the 3 last-mentioned radicals is unsubstituted or substituted by one or more radicals selected from the group consisting of halogen, hydroxyl, cyano, nitro, thiocyanato, (C₁-C₄)-alkoxy, (C₁-C₄)-haloalkoxy, (C₂-C₄)-alkenyloxy, (C₂-C₄)-haloalkenyloxy, (C₁-C₄)-alkylthio, (C₁-C₄)-alkylsulfinyl, (C₁-C₄)-alkylsulfonyl, (C₁-C₄)-haloalkylsulfinyl, (C₁-C₄)-haloalkylsulfonyl and (C₃-C₉)-cycloalkyl, which is unsubstituted or substituted, and phenyl, which is unsubstituted or substituted, and heterocyclyl, which is unsubstituted or substituted, and radicals of the formulae $R'-C(=Z')-$, $R'-C(=Z')-Z-$, $R'-Z-C(=Z')-$, $R'R''N-C(=Z')-$, $R'-Z-C(=Z')-O-$, $R'R''N-C(=Z')-Z-$, $R'-C(=Z')-NR''-$ and $R'R''N-C(=Z')-NR'''-$, in which R', R'' and R''' in each case independently of one another are (C₁-C₆)-alkyl, aryl, aryl-(C₁-C₆)-alkyl, (C₃-C₉)-cycloalkyl or (C₃-C₉)-cycloalkyl-(C₁-C₆)-alkyl, where each of the 5 last-mentioned radicals is unsubstituted or substituted, and in which Z and Z' independently of one another are each an oxygen or sulfur atom,

R^2 is (C₃-C₉)-cycloalkyl, which is unsubstituted or substituted, (C₄-C₉)-cycloalkenyl, which is unsubstituted or substituted, heterocyclyl, which is unsubstituted or substituted, or phenyl, which is unsubstituted or substituted, or

R^3 is hydrogen, (C₁-C₆)-alkyl, aryl or (C₃-C₉)-cycloalkyl, where each of the 3 last-mentioned radicals is unsubstituted or substituted, or a radical of the formula $-N(B^1-D^1)(B^2-D^2)$ or -

- $\text{NR}'\text{-N}(\text{B}^1\text{-D}^1)(\text{B}^2\text{-D}^2)$, in which in each case B^1 , B^2 , D^1 and D^2 are as defined below and
- R' is hydrogen, $(\text{C}_1\text{-C}_6)\text{-alkyl}$ or $[(\text{C}_1\text{-C}_4)\text{-alkyl}]\text{-carbonyl}$,
- R^4 is a radical of the formula $\text{-B}^3\text{-D}^3$, where B^3 and D^3 are as defined below,
- A^1 is straight-chain alkylene having 1 to 5 carbon atoms or straight-chain alkenylene or alkynylene having in each case 2 to 5 carbon atoms, where each of the three last-mentioned diradicals is unsubstituted or substituted by one or more radicals selected from the group consisting of halogen, nitro, cyano, thiocyanato and radicals of the formula $\text{-B}^4\text{-D}^4$, where B^4 and D^4 are as defined below,
- A^2 is a direct bond or straight-chain alkylene having 1 to 4 carbon atoms or straight-chain alkenylene or alkynylene having in each case 2 to 5 carbon atoms, where each of the three last-mentioned diradicals is unsubstituted or substituted by one or more radicals selected from the group consisting of halogen, nitro, cyano, thiocyanato and radicals of the formula $\text{-B}^5\text{-D}^5$, or a divalent radical of the formula V^1 , V^2 , V^3 , V^4 or V^5 ,



where each of the radicals R^6 to R^{27} , in each case independently of one another, is hydrogen, halogen, nitro, cyano, thiocyanato or a radical of the formula $\text{-B}^6\text{-D}^6$,

W^* is in each case an oxygen atom, a sulfur atom or a group of the formula $\text{N}(\text{B}^7\text{-D}^7)$ and B^5 , B^6 , B^7 , D^5 , D^6 and D^7 are as defined below,

B^1 , B^2 , B^3 and B^7 in each case independently of one another are a direct bond or a divalent group of the formulae -C(=Z*)- , $\text{-C(=Z*)}\text{-Z}^{**}\text{-}$, $\text{-C(=Z*)}\text{-NH-}$ or $\text{-C(=Z*)}\text{-NR*}\text{-}$, where Z^* = an oxygen or sulfur atom, Z^{**} = an oxygen or sulfur atom and R^* = $(\text{C}_1\text{-C}_6)\text{-alkyl}$, aryl , $\text{aryl}\text{-(C}_1\text{-C}_6)\text{-alkyl}$, $(\text{C}_3\text{-C}_9)\text{-cycloalkyl}$ or $(\text{C}_3\text{-C}_9)\text{-cycloalkyl}\text{-(C}_1\text{-C}_6)\text{-alkyl}$, where each of the 5 last-mentioned radicals is unsubstituted or substituted,

B^4 , B^5 and B^6 in each case independently of one another are a direct bond or a divalent group of the formulae $-O-$, $-S(O)_p-$, $-S(O)_pO-$, $-O-S(O)_p-$, $-CO-$, $-O-CO-$, $-CO-O-$, $-S-CO-$, $-CO-S-$, $-S-CS-$, $-CS-S-$, $-O-CO-O-$, $-NR^O-$, $-O-NR^O-$, $-NR^O-O-$, $-NR^O-CO-$, $-CO-NR^O-$, $-O-CO-NR^O-$ or $-NR^O-CO-O-$, where p is the integer 0, 1 or 2 and R^O is hydrogen, (C_1-C_6) -alkyl, aryl, aryl- (C_1-C_6) -alkyl, (C_3-C_9) -cycloalkyl or (C_3-C_9) -cycloalkyl- (C_1-C_6) -alkyl, where each of the 5 last-mentioned radicals is unsubstituted or substituted,

D^1 , D^2 , D^3 , D^4 , D^5 and D^6 in each case independently of one another are hydrogen, (C_1-C_6) -alkyl, aryl, aryl- (C_1-C_6) -alkyl, (C_3-C_9) -cycloalkyl or (C_3-C_9) -cycloalkyl- (C_1-C_6) -alkyl, where each of the 5 last-mentioned radicals is unsubstituted or substituted, or in each case two radicals D^5 of two groups $-B^5-D^5$ attached to one carbon atom are attached to one another forming an alkylene group having 2 to 4 carbon atoms which is unsubstituted or substituted by one or more radicals selected from the group consisting of (C_1-C_4) -alkyl and (C_1-C_4) -alkoxy,

$(X)_n$ are n substituents X , where X , in each case independently of one another, is halogen, hydroxyl, amino, nitro, formyl, carboxyl, cyano, thiocyanato, aminocarbonyl or (C_1-C_6) -alkyl, (C_1-C_6) -alkoxy, (C_1-C_6) -alkylthio, mono- (C_1-C_6) -alkylamino, di- (C_1-C_4) -alkylamino, (C_2-C_6) -alkenyl, (C_2-C_6) -alkynyl, $[(C_1-C_6)$ -alkyl]-carbonyl, $[(C_1-C_6)$ -alkoxy]-carbonyl, mono- (C_1-C_6) -alkylamino-carbonyl di- (C_1-C_4) -alkylamino-carbonyl, $N-(C_1-C_6)$ -alkanoyl-amino or $N-(C_1-C_4)$ -alkanoyl- $N-(C_1-C_4)$ -alkylamino, where each of the 13 last-mentioned radicals is unsubstituted or substituted, preferably unsubstituted or substituted by one or more radicals selected from the group consisting of halogen, hydroxyl, amino, nitro, formyl, carboxyl, cyano, thiocyanato, (C_1-C_4) -alkoxy, (C_1-C_4) -haloalkoxy, (C_1-C_4) -alkylthio, (C_1-C_4) -haloalkylthio, mono- (C_1-C_4) -alkylamino, di- (C_1-C_4) -alkylamino, (C_3-C_9) -cycloalkyl, (C_3-C_9) -cycloalkyl-amino, $[(C_1-C_4)$ -alkyl]-carbonyl, $[(C_1-C_4)$ -alkoxy]-carbonyl, aminocarbonyl, mono- (C_1-C_4) -alkylamino-carbonyl, di- (C_1-C_4) -alkylamino-carbonyl, phenyl, phenoxy, phenylthio, phenylcarbonyl, heterocyclyl, heterocycloxy, heterocyclthio and heterocyclylamino, where each of the 8 last-mentioned radicals is unsubstituted or substituted by one or more substituents selected from the group consisting of halogen, nitro, cyano, (C_1-C_4) -alkyl, (C_1-C_4) -alkoxy, (C_1-C_4) -alkylthio, (C_1-C_4) -haloalkyl, (C_1-C_4) -haloalkoxy, formyl, (C_1-C_4) -alkyl-carbonyl and (C_1-C_4) -alkoxy-carbonyl,

or (C₃-C₉)-cycloalkyl, (C₃-C₉)-cycloalkoxy, (C₃-C₉)-cycloalkylamino, phenyl, phenoxy, phenylthio, phenylcarbonyl, heterocyclyl, heterocyclyloxy, heterocyclylthio or heterocyclylamino, where each of the 11 last-mentioned radicals is unsubstituted or substituted, or two adjacent radicals X together are a fused-on cycle having 4 to 6 ring atoms which is carbocyclic or contains hetero ring atoms selected from the group consisting of O, S and N and which is unsubstituted or substituted by one or more radicals selected from the group consisting of halogen, (C₁-C₄)-alkyl and oxo,


n is 0, 1, 3, 4 or 5 and

Heterocyclyl in the radicals mentioned above, independently of one another, is in each case a heterocyclic radical having 3 to 7 ring atoms and 1 to 3 heteroatoms selected from the group consisting of N, O and S,

where

- a) the total of the carbon atoms in the radicals A¹ and A²-R² is at least 6 carbon atoms or
- b) the total of the carbon atoms in the radicals A¹ and A²-R² is 5 carbon atoms and A¹ = a group of the formula -CH₂- or -CH₂CH₂- and R¹ = (C₁-C₄)-alkyl, (C₁-C₄)-haloalkyl, (C₂-C₆)-haloalkenyl or (C₃-C₉)-cycloalkyl, which is unsubstituted or substituted;

- compounds of the formula (IV) or their salts



in which

R¹ and R² in each case independently of one another are hydrogen, amino, alkylamino or dialkylamino having in each case 1 to 6 carbon atoms in the alkyl radical, an acyclic or cyclic hydrocarbon radical or hydrocarbonoxy radical having in each case 1 to 10 carbon atoms or a

heterocyclyl radical, heterocyclyloxy radical, heterocyclylthio radical or heterocyclylamino radical having in each case 3 to 6 ring atoms and 1 to 3 hetero ring atoms selected from the group consisting of N, O and S, where each of the five last-mentioned radicals is unsubstituted or substituted, or an acyl radical or

R^1 and R^2 together with the nitrogen atom of the group NR^1R^2 are a heterocyclic radical having 3 to 6 ring atoms and 1 to 4 hetero ring atoms, where any further hetero ring atoms present in addition to the nitrogen atom are selected from the group consisting of N, O and S and the radical is unsubstituted or substituted,

R^3 is halogen, cyano, thiocyanato, nitro or a radical of the formula $-Z^1-R^7$,

R^4 is hydrogen, amino, alkylamino or dialkylamino having in each case 1 to 6 carbon atoms in the alkyl radical, an acyclic or cyclic hydrocarbon radical or hydrocarbonoxy radical having in each case 1 to 10 carbon atoms or a heterocyclyl radical, heterocyclyloxy radical or heterocyclylamino radical having in each case 3 to 6 ring atoms and 1 to 3 hetero ring atoms selected from the group consisting of N, O and S, where each of the five last-mentioned radicals is unsubstituted or substituted, or an acyl radical,

R^5 is halogen, cyano, thiocyanato, nitro or a radical of the formula $-Z^2-R^8$,

R^6 , in the case where $n=1$, or the radicals R^6 in each case independently of one another, if n is greater than 1, is/are halogen, cyano, thiocyanato, nitro or a group of the formula $-Z^3-R^9$,

R^7 , R^8 , R^9 in each case independently of one another are

- hydrogen or
- an acyclic hydrocarbon radical, where carbon atoms in the chain may be substituted by heteroatoms selected from the group consisting of N, O and S, or
- a cyclic hydrocarbon radical or
- a heterocyclic radical,

where each of the 3 last-mentioned radicals is unsubstituted or substituted,

Z^1, Z^2, Z^3 in each case independently of one another are

- a direct bond or
- a divalent group of the formula $-O-$, $-S(O)_p-$, $-S(O)_p-O-$, $-O-S(O)_p-$, $-CO-$, $-CS-$, $-S-CO-$, $-CO-S-$, $-O-CS-$, $-CS-O-$, $-S-CS-$, $-CS-S-$, $-OCO-$, $-CO-O-$, $-NR'-$, $-O-NR'-$, $-NR'-O-$, $-NR'-CO-$ or $-CO-NR'-$, where

$p = 0, 1$ or 2 and R' is hydrogen, alkyl having 1 to 6 carbon atoms, phenyl, benzyl, cycloalkyl having 3 to 6 carbon atoms or alkanoyl having 1 to 6 carbon atoms,

Y^1, Y^2, Y^3 and, if m is 2, 3 or 4, further groups Y^2 are, in each case independently of one another,


- a divalent group of the formula CR^aR^b , where R^a and R^b are identical or different and are in each case a radical selected from the group of the radicals possible for R^7 to R^9 , or
- a divalent group of the formula $-O-$, $-CO-$, $-C(=NR^*)-$, $-S(O)_q-$, $-NR^*-$ or $-N(O)-$, where $q = 0, 1$ or 2 and R^* is hydrogen or alkyl having 1 to 4 carbon atoms, or
- Y^1 or Y^3 is a direct bond,

where two oxygen atoms of groups Y^2 and Y^3 are not adjacent,

m is 1, 2, 3 or 4,

n is 0, 1, 2, 3 or 4;

- substituted 2,4-diamino-1,3,5-triazines of the formula (V),



in which

- R¹ is hydrogen or unsubstituted or hydroxyl-, cyano-, halogen- or C₁-C₄-alkoxy-substituted alkyl having 1 to 6 carbon atoms,
- R² is hydrogen, formyl, in each case unsubstituted or cyano-, halogen- or C₁-C₄-alkoxy-substituted alkyl, alkylcarbonyl, alkoxy carbonyl or alkylsulfonyl having in each case 1 to 6 carbon atoms in the alkyl groups, or is unsubstituted or cyano-, halo-C₁-C₄-alkyl-, C₁-C₄-alkoxy-, halo-C₁-C₄-alkoxy- or C₁-C₄-alkoxy-carbonyl-substituted phenylcarbonyl, naphthylcarbonyl, phenylsulfonyl or naphthylsulfonyl,
- R³ is unsubstituted or cyano-, halogen- or C₁-C₄-alkoxy-substituted alkyl having 1 to 6 carbon atoms or is unsubstituted or cyano-, halogen- or C₁-C₄-alkyl-substituted cycloalkyl having 3 to 6 carbon atoms,
- X is a substituent selected from the group below:
hydroxyl, cyano, nitro, halogen, in each case unsubstituted or hydroxyl-, cyano- or halogen-substituted alkyl or alkoxy having in each case 1 to 6 carbon atoms, in each case unsubstituted or halogen-substituted alkylcarbonyl, alkoxy carbonyl, alkylthio, alkylsulfinyl or alkylsulfonyl having in each case 1 to 6 carbon atoms in the alkyl groups, in each case unsubstituted or hydroxyl-, cyano-, nitro-, halogen-, C₁-C₄-alkyl, C₁-C₄-haloalkyl-, C₁-C₄-alkoxy or C₁-C₄-haloalkoxy-substituted phenyl or phenoxy, and
- Z is hydrogen, hydroxyl, halogen, in each case unsubstituted or hydroxyl-, cyano-, nitro-, halogen-, C₁-C₄-alkoxy-, C₁-C₄-alkyl-carbonyl-, C₁-C₄-alkoxy-carbonyl-, C₁-C₄-alkylthio-, C₁-C₄-alkylsulfinyl- or C₁-C₄-alkylsulfonyl-substituted alkyl, alkoxy, alkylcarbonyl, alkoxy carbonyl, alkylthio, alkylsulfinyl or alkylsulfonyl, having in each case 1 to 6 carbon atoms in the alkyl groups, in each case unsubstituted or halogen-substituted alkenyl or alkynyl having in each case 2 to 6 carbon atoms or is unsubstituted or cyano-, halogen- or C₁-C₄-alkyl-substituted cycloalkyl having 3 to 6 carbon atoms.

and

- (B) is one or more herbicides selected from the group of compounds consisting of
- (B1) foliar- and/or soil-acting herbicides which are active against monocotyledonous harmful plants selected from the group consisting of

- (B1.1.1) isoproturon,
(B1.1.2) chlorotoluron,

- (B1.2.1) flufenacet,
 - (B1.2.2) pendimethalin,
 - (B1.2.3) prosulfocarb,
 - (B1.3.1) clodinafop-propargyl,
 - (B1.3.2) diclofop-methyl,
 - (B1.3.3) fenoxaprop-P-ethyl and fenoxaprop-ethyl,
 - (B1.3.4) quizalofop-P and its salts and esters and quizalofop and its salts
- and esters,
- (B1.3.5) fluazifop-P and its esters and fluazifop and its esters,
 - (B1.3.6) haloxyfop and haloxyfop-P and their esters,
 - (B1.3.7) propaquizafop (PM, p. 1021-1022),
 - (B1.3.8) cyhalofop and its esters,
 - (B1.4.1) sethoxydim,
 - (B1.4.2) cycloxydim
 - (B1.4.3) clethodim,
 - (B1.4.4) clefoxidim,
 - (B1.4.5) tralkoxidim,
 - (B1.5.1) dimethenamid,
 - (B1.5.2) penthoxamid,
 - (B1.5.3) butachlor,
 - (B1.5.4) pretilachlor,
 - (B1.6.1) imazamethabenz-methyl
 - (B1.6.2) simazin
 - (B1.6.3) molinate
 - (B1.6.4) thiobencarb
 - (B1.6.4) MY 100,
 - (B1.6.5) anilofos,
 - (B1.6.6) cafenstrole,
 - (B1.6.7) mefenacet,
 - (B1.6.8) fentrazamid,
 - (B1.6.9) thiazopyr,

- (B1.6.10) oxadiazon,
- (B1.6.11) esprocarb,
- (B1.6.12) pyributicarb,
- (B1.6.13) azimsulfuron,
- (B1.6.14) AEB391 and related azoles,
- (B1.6.15) thenylchlor,
- (B1.6.16) pentoxazone,
- (B1.6.17) pyriminobac and pyriminobac-methyl,
- (B1.6.18) flucarbazone and its salts and
- (B1.6.19) procarbazon and its salts,

(B2) herbicides which are active predominantly against dicotyledonous harmful plants selected from the group consisting of

- (B2.1.1) tribenuron-methyl,
- (B2.1.2) thifensulfuron and its esters,
- (B2.1.3) prosulfuron,
- (B2.1.4) amidosulfuron,
- (B2.1.5) chlorimuron and its esters,
- (B2.1.6) halosulfuron and its esters and salts,
- (B2.1.7) LAB271272, (= tritosulfuron),
- (B2.1.8) bensulfuron-methyl,
- (B2.1.9) ethoxysulfuron,
- (B2.1.10) cinosulfuron,
- (B2.1.11) pyrazosulfuron and its esters,
- (B2.1.12) imazosulfuron,
- (B2.1.13) cyclosulfamuron,
- (B2.2.1) MCPA,
- (B2.2.2) 2,4-D,
- (B2.2.3) dichlorprop,
- (B2.2.4) mecoprop-(P),
- (B2.2.5) fluoroxypyr,
- (B2.2.6) dicamba,

- (B2.2.7) clopyralid,
- (B2.2.8) picloram,
- (B.2.3.1) bromoxynil,
- (B.2.3.2) ioxynil,
- (B2.4.1) fluoroglycofen-ethyl,
- (B2.4.2) aclonifen,
- (B2.4.3) acifluorfen and its salts,
- (B2.5.1) cloransulam and its esters
- (B2.5.2) florasulam,
- (B2.6.1) bentazone,
- (B2.6.2) bifenox,
- (B2.6.3) carfentrazone-ethyl,
- (B2.6.4) pyraflufen,
- (B2.6.5) pyridate,
- (B2.6.6) linuron,
- (B2.6.7) diflufenzopyr and its salts,
- (B2.6.8) cinidon-ethyl,
- (B2.6.9) clopyralid and its salts and esters,
- (B2.6.10) metribuzin,
- (B2.6.11) picolinafen,
- (B2.6.12) clomazone,
- (B2.6.13) bromobutide,
- (B2.6.14) benfuresate,
- (B2.6.15) dithiopyr and
- (B2.6.16) triclopyr and its salts and esters,

(B3) herbicides which are active against monocotyledonous and dicotyledonous harmful plants selected from the group consisting of

- (B3.1.1) metsulfuron and its esters,
- (B3.1.2) triasulfuron,
- (B3.1.3) chlorsulfuron,
- (B3.1.4) iodosulfuron-methyl,

(B3.1.5)	AEF060,
(B3.1.6)	sulfosulfuron,
(B3.1.7)	flupyrsulfuron and its salts,
(B3.1.8)	nicosulfuron,
(B3.1.9)	rimsulfuron,
(B3.1.10)	primisulfuron and esters,
(B3.1.11)	AEF360,
(B3.2.1)	cyanazin
(B3.2.2)	atrazin
(B3.2.3)	terbuthylazin,
(B3.2.4)	terbutryn,
(B3.3.1)	acetochlor
(B3.3.2)	metolachlor,
(B3.3.3)	alachlor,
(B3.4.1)	clomazone,
(B3.4.2)	diflufenican,
(B3.4.3)	flumetsulam,
(B3.4.4)	flurtamone,
(B3.4.5)	isoxaflutole,
(B3.4.6)	metosulam,
(B3.4.7)	metribuzin,
(B3.4.8)	paraquat (salts),
(B3.4.9)	benoxacor,
(B3.4.10)	sulcotrione,
(B3.4.11)	mesotrione,
(B3.4.12)	quinclorac,
(B3.4.13)	propanil,
(B3.4.14)	bispyribac, bispyribac-Na,
(B3.4.15)	LGC 40863 (pyribenzoxim),
(B3.4.16)	oxadiargyl,
(B3.4.17)	norflurazon,

- (B3.4.18) fluometuron,
- (B3.4.19) methylarsonic acid and its salts (DSMA, MSMA).
- (B3.4.20) prometryn,
- (B3.4.21) trifluralin,

(B4) herbicides which are active against monocotyledonous and dicotyledonous harmful plants and which can be employed specifically in tolerant crops or on non-crop land, selected from the group consisting of

- (B4.1.1) glufosinate,
- (B4.1.2) glufosinate monoammonium salt,
- (B4.1.3) L-glufosinate,
- (B4.1.4) L-glufosinate monoammonium salt,
- (B4.1.5) bilanafos,
- (B4.2.1) glyphosate,
- (B4.2.2) glyphosate monoisopropylammonium salt,
- (B4.2.3) glyphosate sodium salt,
- (B4.2.4) sulfosate,
- (B4.3.1) imazapyr,
- (B4.3.2) imazethapyr
- (B4.3.3) imazamethabenz, and its salts and esters,
- (B4.3.4) imazamox and its salts and esters,
- (B4.3.5) imazaquin and its salts and esters,
- (B4.3.6) imazapic (AC 263,222) and its salts and esters
- (B4.4.1) WC9717 or CGA276854,
- (B4.4.2) azafenidin,
- (B4.4.3) diuron and
- (B4.4.4) oxyfluorfen,

and, if appropriate, their agriculturally useful salts.

20. (Previously presented) The herbicide combination as claimed in claim 19, wherein the one or more herbicidally active aminotriazine compound (A) is selected from the group consisting of:

(A1) 4-amino-6-(1-fluoro-1-methylethyl)-2-(3-phenyl-1-cyclobutylamino)-1,3,5-triazine;
and

wherein the one or more herbicides (B) is selected from the group consisting of:

- (B1.1.1) isoproturon;
- (B1.2.1) Flufenacet;
- (B1.2.2) Pendamethalin;
- (B1.3.3) Fenoxaprop-P-ethyl
- (B.1.6.4) MY100;
- (B1.6.14) AEB391;
- (B2.1.1) Tribenuron-methyl;
- (B2.1.4) Amidosulfuron;
- (B2.2.4) Mecoprop-P;
- (B2.3.1) Bromoxynil;
- (B2.3.2) Ioxynil;
- (B2.5.2) Florsulam;
- (B2.6.3) Carfentrazone-ethyl;
- (B3.1.11) AEF360 with isoxadifen-ethyl; and
- (B3.4.2) Duflufenican;

21. (Previously presented) The herbicide combination as claimed in claim 19, which comprises, as component (A), one or more triazine derivatives of the formulae (A1), (A2) and (A3):

